**Research Article** 

# Characterization of *Citrus aurantifolia* 'Nimas Agrihorti' Cultivar as a Potential Biopharmaceutical Commodity and Its Economic Feasibility in Indonesia

# Emi Budiyati<sup>1\*</sup>, Budi Waluyo<sup>2</sup>, Moch. Dawam Maghfoer<sup>2</sup>, Mia Kosmiatin<sup>1</sup>, Anang Triwiratno<sup>1</sup>, Zainuri Hanif<sup>5</sup>, Anis Andrini<sup>3</sup>, Faridha Hanim<sup>2</sup>, & Siti Agus Tina<sup>4</sup>

<sup>1</sup>Research Center for Horticulture, National Agency for Research and Innovation (BRIN), Cibinong, Bogor 16912, West Java, Indonesia

<sup>2</sup> Faculty of Agriculture, Brawijaya University, Malang 65145, East Java, Indonesia

<sup>3</sup> Agricultural Standarization and Instruments Agency, Ministry of Agriculture, Jakarta 12540, Indonesia

<sup>4</sup> Faculty of Agriculture, IPB University, IPB Dramaga Campus, Bogor 16680, West Java, Indonesia

<sup>5</sup>Research Center for Behavioral and Circular Economics, National Research and Innovation Agency (BRIN), Jakarta, Indonesia

\*Corresponding author: emib140666@gmail.com

#### ABSTRACT

'Nimas Agrihorti' is a citrus cultivar developed to fulfil the increasing domestic demand for biopharmaceutical-grade citrus in Indonesia. Its development aligns with the growing public interest in health-promoting agricultural products. This study aimed to evaluate the morphological characteristics, chemical composition, and economic viability of 'Nimas Agrihorti' as a biopharmaceutical citrus. Field observations were conducted from June 2019 to October 2020 at the Banjarsari Experimental Field, Bayeman Village, Probolinggo, East Java. Morphological assessments followed the Descriptor List for Citrus by IPGRI, and chemical analyses were performed at the Post-Harvest Laboratory of Brawijaya University, Malang. Economic feasibility was assessed by calculating production costs and revenues over a 2-year cultivation cycle. 'Nimas Agrihorti' exhibited favorable morphological traits, including large fruit size (72-82 g), yellow skin, and sweet taste. Chemical analysis revealed vitamin C content of 34.8 mg/100 g and low acidity at 0.45%. The total production cost over 2 years was IDR 42,640,000.00 (equivalent to USD 2,584), with total revenue amounting to IDR 77,360,000.00 (equivalent to USD 4,688), resulting in a Revenue-Cost (R/C) ratio of 2.81. The combination of desirable morphological and chemical characteristics, along with strong economic returns, indicates that 'Nimas Agrihorti' holds significant potential for development as a biopharmaceutical citrus cultivar in Indonesia.

Keywords: citrus, cultivar, biopharmaceutical products, economic feasibility

#### Introduction

The availability of plant-based biopharmaceutical products for medicinal use in Indonesia has declined over the past few decades. According to BRIN-Indonesia National Research and Innovation Agency (2023), Indonesia remained behind in producing sufficient raw materials for biopharmaceutical drugs, which possessed therapeutic benefits, particularly for enhancing immune function (Annisa et al., 2022). These benefits could be obtained through direct human consumption or via industrial extraction processes (Fahrurin et al., 2023). However, data from BPS-Statistics Indonesia (2024) indicated a positive trend for citrus as a biopharmaceutical commodity showing increased production in 2021 and 2022, reaching 13,103,964 kg and 25,175,958 kg, respectively.

Citrus fruits are widely appreciated not only for their pleasant flavour but also for their abundance of bioactive compounds and essential nutrients. They are a significant source of phenolic constituents, flavonoids, phenolic including acids. terpenoids, coumarins. limonoids. carotenoids, and pectin (Budiarto et al., 2024). Additionally, citrus is nutritionally rich in compounds such as ascorbic acid (vitamin C), tocopherols and tocotrienols (vitamin E), as well as trace minerals like selenium, zinc. copper, iron, and manganese (Saini et al., 2022). The peel of fruits contains essential citrus oils, carotenoids, flavonoids, and other bioactive molecules known for their potent antioxidant properties. Moreover, citrus valuable seeds offer nutritional components, including essential proteins, ascorbic acid, fatty acids, phytosterols, tocopherols, limonoids, dietary fiber, and flavonoids (Saini et al., 2022).

Citrus is recognized as one of the fruit sources of carotenoids, major compounds known for their potential in cancer prevention. The concentration of carotenoids in citrus fruits can vary depending environmental on and cultivation conditions (Gebregziabher et al., 2023). Citrus extracts are particularly in flavanones. with hesperidin rich identified as the most abundant phenolic compound (Khan et al., 2014; Kumar et al., 2021; Mitra et al., 2022). Phenolic compounds derived from citrus exhibit strong biological activities, including anticarcinogenic, antimutagenic (de Luna et al., 2023), and antitumor properties (Mueed et al., 2023). They have been associated with reducing the risk of colorectal (Gao et

al., 2022), esophageal, gastric, and stomach cancers (Tullio et al., 2020), as well as contributing to the prevention of stroke, cholesterol regulation, and exhibiting antioxidant effects (Alageel, 2023; Hijaz et al., 2020). Furthermore, these compounds possess anti-inflammatory (Denaro et al., 2021), antifungal, and antithrombotic activities. Hesperidin, in particular, has demonstrated potential in combating coronavirus infections. lowering cholesterol levels, and enhancing bile acid secretion (Saini et al., 2022). Regular consumption of citrus fruits is thus highly beneficial to human health, supported by their rich nutrient profile and palatable taste (Liu et al., 2022).

Extensive research has been conducted on the health benefits of citrus species. For instance, Shafreen et al. (2018) investigated the interaction between limederived phenolic compounds and human serum proteins. Shimada et al. (2022) reported the bioactivity of kabosu (Citrus sphaerocarpa) in inhibiting angiogenesis and the migration of breast cancer cells. Similarly, Anuchapreeda et al. (2020) highlighted the use of kaffir lime leaves (*Citrus hystrix*) both as a culinary ingredient and in traditional medicine. Additional studies by Prommaban and Chaiyana (2022) demonstrated the antiaging and anti-melanogenic properties of essential oils extracted from the leaves and peels of *Citrus aurantifolia* (Christm.) Swing and Citrus reticulata Blanco cv. Shogun. While numerous studies have explored the bioactive components and therapeutic effects of citrus fruits, relatively few have focused on the development of new cultivars. One such example is the 'Nimas Agrihorti' cultivar, which offers health benefits-particularly for low-fat applications dietary and diabetes management-despite its naturally sweet flavor profile.

The development of new citrus cultivars is anticipated to support initiatives by the Directorate General of Horticulture, Ministry of Agriculture, aimed at reducing the influx of imported citrus fruits, which are becoming increasingly accessible in Indonesia (Rahayu et al., 2020). Achieving this goal requires efforts to enhance fruit quality, ensure year-round availability, and increase production volume. One approach involves the cultivation of the 'Nimas Agrihorti' citrus cultivar (Figure 1). As a newly introduced cultivar, 'Nimas Agrihorti' citrus developed in 2012 by the Citrus and Subtropical Fruit Research Center has been identified as possessing high nutritional value and the potential for dual use—as a fresh consumption product and as a raw material for biopharmaceutical applications. In addition to its nutritional qualities, 'Nimas Agrihorti' also exhibits favorable productivity and economic value. the advantages of Besides 'Nimas Agrihorti', this study was conducted to evaluate the potential of 'Nimas Agrihorti' as a biopharmaceutical citrus by analyzing its chemical composition and assessing its economic feasibility.



**Fig. 1.** *Citrus aurantifolia* 'Nimas Agrihorti' fruits in the field. Source: research documentation (2015).

#### **Materials and Methods**

The study was conducted on 'Nimas Agrihorti' mother plants aged 2.5 to 3.5 years, with two harvest cycles, grown in an experimental plot of approximately 1,000 m<sup>2</sup>. Field observations were conducted at the Banjarsari Experimental Field, Bayeman Village, Probolinggo, East Java. The site was characterized by alluvial soil, an elevation of 1 m above sea level, an average annual rainfall of 1,200 mm, approximately 90 rainy days per year, a temperature range of 21-28°C, relative humidity of 76%, and soil pH between 6 and 7. Observations and plant identification were carried out from June 2019 to October 2020. Morphological characteristics were evaluated using the Descriptor List for Citrus developed by International Plant Genetic Resources Institute (IPGRI), while analyses of the fruit chemical properties were conducted at the Post-Harvest Laboratory, Brawijaya University, Malang.

# Variables of Observation

The qualitative traits assessed in this study included overall plant morphology, such as stem, leaf, and fruit shape and color. shape was evaluated through Stem transverse stem sections, while stem color was visually assessed and confirmed through consensus by five evaluators. Fruit skin color, along with the color of the flesh, leaves, and flowers, was determined by comparing samples to a standardized color Quantitative chart. characteristics encompassed parameters such as plant height, yield (number and weight of fruits), fruit dimensions, vitamin content, stem diameter, leaf size, number of fruit segments, flower count, and fruit count. Additional measurements included total fruit per plant, average fruit weight, fruit length and diameter, sugar content, and results from chemical composition analysis.

# Variable of Farm Feasibility

According to Pebrianto et al. (2023), farm feasibility is evaluated using the Revenue-Cost (R/C) ratio. A ratio equal to 1 indicates a break-even point, meaning the capital investment is fully recovered. If the R/C ratio exceeds 1, the farming activity is considered profitable, whereas a ratio below 1 reflects a financial loss.

## **Results and Discussion**

Characteristics of 'Nimas Agrihorti'

'Nimas Agrihorti' was obtained through hybridization between citrus and sweet orange, resulting in distinctive traits such as a plant height ranging from 156 to 214 cm, a shrub-like canopy structure, and stem diameters of 4–6 cm with a greyish coloration. Unlike typical lemon plants, which generally begin fruit production at heights of 3–5 m, 'Nimas Agrihorti' demonstrates precocious fruiting ability at significantly lower plant heights, indicating its early-yielding advantage.

The hybridization between citrus and sweet orange has resulted in the morphological superiority of the 'Nimas Agrihorti' cultivar, particularly in terms of flavor profile, fruit size, and other agronomic traits. The leaf shape of 'Nimas Agrihorti' exhibited an oval shape, with lengths ranging from 7 to 10 cm and widths between 4 and 6 cm. The plant produced compound flowers. Illustrations of the leaf and flower morphology of 'Nimas Agrihorti' are presented in Figure 2.



**Fig. 2.** Flowers (a) dan leaves (b) of *Citrus aurantifolia* 'Nimas Agrihorti'.

The leaf and flower morphology of 'Nimas Agrihorti' resembled that of common citrus species, displaying an oval to elliptical leaf shape (Ernawati et al., 2023). Based on the findings of Adlini et al. (2020), citrus leaves typically ranged from 5.5–13 cm in length and 2–7.5 cm in width, while the Cancer Chemoprevention Research Center reported leaf dimensions of 2.5–9 cm in length and 2–5 cm in width (CCRC UGM, 2008). Compared to these references, the leaf size of 'Nimas Agrihorti' falls within the broader range, classifying it in the large-leaf category. This characteristic is important, as leaf size is known to influence the photosynthetic rate, which is associated with plant productivity (Setyanti et al., 2013; Yustiningsih, 2019; Zulkifli et al., 2022). When compared to other lime cultivars, such as 'Nipis Borneo', 'Nimas Agrihorti' demonstrated superior morphological traits, as summarized in Table 1.

Table 1. Comparative morphologicalcharacteristics of Citrus aurantifolia'Nimas Agrihorti' and 'Nipis Borneo'cultivars.

| Plant                             | 'Nimas          | 'Nipis Borneo'                        |  |
|-----------------------------------|-----------------|---------------------------------------|--|
| morphology                        | Agrihorti'      |                                       |  |
| Canopy shape                      | Erect           | Round                                 |  |
| Stem color                        | Grey            | Greenish brown                        |  |
| Leaf shape                        | Ovate           | Elliptic                              |  |
| Leaf color                        | Green           | Green                                 |  |
| Flower shape                      | Compound        | Compound                              |  |
| Flower color                      | Green           | Light green                           |  |
| <ul> <li>Flower petals</li> </ul> | White           | White                                 |  |
| <ul> <li>Stamen</li> </ul>        | Yellow          | Dark yellow                           |  |
| <ul> <li>Another</li> </ul>       | Yellowish green | Light yellow                          |  |
| Flower type                       | Compound        | Compound                              |  |
| Flowering time                    | Continuously    | January, March, July,<br>and November |  |
| Harvest time                      | Continuously    | April, June,<br>September, and        |  |
|                                   |                 | December                              |  |
| Fruit shape                       | Ellipsoid       | Round                                 |  |
| Skin color                        | Yellow          | Yellow                                |  |
| Flesh color                       | White           | White                                 |  |
| Flesh taste                       | Sweet           | Sour                                  |  |
| Seed shape                        | Ovoid           | Elliptic                              |  |
| Seed color                        | Cream           | White bone                            |  |

Although both 'Nipis Borneo' and 'Nimas Agrihorti' cultivars shared similar exocarp and pulp coloration, 'Nimas Agrihorti' was notably sweeter in taste. In addition to differences in morphological traits of the tree and fruit, a comparative analysis of their quantitative attributes is presented in Table 2.

The quantitative character indicated that 'Nimas Agrihorti' was a citrus cultivar characterized by high water content (60%) and low acidity (0.45%) (Table 2). Although its sugar content (8.3%) was lower than that of 'Nipis Borneo' (10.8%), the overall taste profile of 'Nimas remained favorable. Agrihorti' These attributes contribute to its superior sensory qualities compared to other citrus cultivars. Given these compositional advantages, 'Nimas Agrihorti' demonstrated promising potential for development as а biopharmaceutical or medicinal citrus cultivar.

**Table 2.** Quantitative characterization of *Citrus aurantifolia* 'Nimas Agrihorti' in 2020 and comparative analysis with 'Nipis Borneo'.

| Characteristics                   | 'Nimas         | 'Nipis         |
|-----------------------------------|----------------|----------------|
|                                   | Agrihorti'     | Borneo'        |
| Plant height (cm)                 | 156-214        | 550            |
| Stem diameter (cm)                | 6.51           | 13.8-29.0      |
| Leaf size                         |                |                |
| <ul> <li>Length (cm)</li> </ul>   | 11.48          | 6.3-8.0        |
| <ul> <li>Width (cm)</li> </ul>    | 5.46           | 2.8 - 4.1      |
| Fruit size                        |                |                |
| <ul> <li>Length (cm)</li> </ul>   | 5-6            | 3.5-4.7        |
| <ul> <li>Diameter (cm)</li> </ul> | 5.65-5.92      | 3.7-5.1        |
| Thickness of fruit skin           | 2.40-3.81      | 1-2            |
| (mm)                              |                |                |
| Water content (%)                 | 60.2           | 37.25-43.96    |
| Sugar content (%)                 | 8.3            | 10.2 - 10.8    |
| Vitamin C content                 | 34.8           | 53.66-55.58    |
| (mg/100 g)                        |                |                |
| Acid content (%)                  | 0.45           | 11.86-15.53    |
| Number of segments                | 10             | 10-12          |
| per fruit                         |                |                |
| Weight per fruit (g)              | 72-80          | 30-70          |
| Fruit number per plant            | 80-120 (on     | 3,000-4,000    |
|                                   | pot)           | (in the field) |
| Percentage of edible              | 70–75          | 71–74          |
| fruit part (%)                    |                |                |
| Production per plant              | 10-15 (on pot) | 220,223 (in    |
| (kg)                              |                | the field)     |
| Age of mother plant               | 11             | 25             |
| (years)                           |                |                |

Potentials 'Nimas Agrihorti' as Biopharmaceutical

The fruit production of 'Nimas Agrihorti' began at 2.5 to 3 years after grafting, with each plant yielding approximately 10 kg of fruit. With a planting density of 400-500 trees/ha, the estimated yield reached around 5,000 kg/ha. As the plant matures, its productivity tended to increase, 'Nimas Agrihorti' also showed strong potential for commercial development due to its competitive traits, including large fruit size (72-82 g), yellow exocarp, sweet flavor, a vitamin C content of 34.8 mg/100 g, and low acidity at 0.45%. A summary of its chemical properties relevant to biopharmaceutical development is presented in Table 3.

The extract yield of 'Nimas Agrihorti' was 44.13%, placing it in the moderate category relative to other citrus cultivars. The extract yield was lower than that of 'Sitaya Agrihorti' (59.05%) and 'Krisma Agrihorti' (54.67%), while it was notably higher than 'Nagamik' (14.66%) and 'Sari Agrihorti' (22.88%). This intermediate extract content indicates its suitability as a raw material for citrus-based beverages with competitive functional properties. The sugar content of 'Nimas Agrihorti' was 8.30%, which is comparable 'Krisma Agrihorti' (8.68%) to and 'Chokun' (8.60%), and significantly higher than that of 'Sitaya Agrihorti' (4.21%) and 'Sari Agrihorti' (3.11%). The natural sweetness of 'Nimas Agrihorti' enhanced its appeal to consumers seeking flavorful, health-oriented vet fruit options, positioning it well for fresh consumption or use in functional beverage formulations.

**Table 3.** Chemical composition analysis of *Citrus aurantifolia* 'Nimas Agrihorti' and other citrus cultivars.

| Cultivars  | Extract<br>(%) | Sugar<br>(%) | Acid<br>(%) | Vitamin C<br>(mg/100 |
|------------|----------------|--------------|-------------|----------------------|
|            |                |              |             | g)                   |
| 'Krisma    | 54.67          | 8.68         | 1.85        | 22.52                |
| Agrihorti  |                |              |             |                      |
| 'Chokun'   | 36.16          | 8.60         | 0.50        | 17.98                |
| 'Sitaya    | 59.05          | 4.21         | 0.24        | 21.55                |
| Agrihorti' |                |              |             |                      |
| 'Nagamik'  | 14.66          | 5.31         | 13.70       | 26.17                |
| 'Sari      | 22.88          | 3.11         | 19.16       | 31.46                |
| Agrihorti' |                |              |             |                      |
| 'Nimas     | 44.13          | 8.30         | 0.45        | 33.47                |
| Agrihorti' |                |              |             |                      |
| 'Nipis     | -              | 8.60         | 3.392       | 34.80                |
| Borneo'    |                |              |             |                      |

One of the distinguishing features of 'Nimas Agrihorti' was its exceptionally low acidity level at 0.45%, the lowest among the citrus cultivars evaluated. This significantly differentiates it from high-acid cultivars such as 'Nagamik' (13.70%) and 'Sari Agrihorti' (19.16%). The mild acidity enhances its palatability and makes it suitable for individuals with dietary limitations, such as those experiencing acid reflux or gastric sensitivity. Furthermore, the low acid profile expands its potential for use in culinary and industrial products that require low-acid extracts.

In terms of nutritional composition, 'Nimas Agrihorti' had high vitamin C content of 33.47 mg/100 g, closely following 'Nipis Borneo' (34.80 mg/100 g) and exceeding other cultivar such as 'Krisma Agrihorti' (22.52 mg/100 g) and 'Sitaya Agrihorti' (21.55 mg/100 g). This substantial vitamin С concentration supports its use as a functional food with antioxidant properties, contributing to immune function, reduction of oxidative and and collagen carnitine stress. biosynthesis-factors critical to maintaining overall health and body resistance (Ivakdalam & Rehena, 2020; Yulianto et al., 2022). The increasing global consumption of citrus extracts reflects rising awareness of the health benefits associated with their bioactive components (Ali Abadi et al., 2022; Shafiee & Minaei, 2018).

'Nimas Agrihorti' also demonstrated a rich profile of phenolic compounds, particularly unique flavonoids such as methoxyflavones (MFs), coumarins (Cs), and furanocoumarins (FCs), known to enhance antioxidant activity (Li et al., 2021; da Silva et al., 2013). The extract is especially abundant in hesperidin-a flavanone-followed dominant by narirutin, commonly found in citrus species. These phenolic compounds were reported to exhibit potent bioactivity, anticarcinogenic, including antitumor. antimutagenic effects, and the ability to lower risks associated with colorectal, esophageal, gastric, and stomach cancers, as well as stroke (Gao et al., 2022).

#### Economic Potential Based on Feasibility of 'Nimas Agrihorti' Cultivation

The assessment of farm feasibility for newly developed citrus cultivars offers valuable insights into their economic viability and sustainability. 'Nimas Agrihorti' presents strong commercial potential due to its favorable combination of sweetness and elevated vitamin C content. These traits make it suitable for fresh consumption, and it is also wellpositioned for use in functional food, pharmaceutical, and nutraceutical product development. As a result, it holds promise as a high-value commodity and a multifunctional ingredient across various industries.

The morphological characteristics of 'Nimas Agrihorti', as described in section 3.1, demonstrated superiority compared to other citrus types. The introduction and promotion of new cultivars were also assessed through their economic viability, which serves as a key consideration for growers in determining the sustainability of continued cultivation. The economic performance of 'Nimas Agrihorti' was analyzed in the following section.

The cultivation of the 'Nimas Agrihorti' citrus cultivar demonstrated strong economic feasibility, with a total production cost of IDR 42,640,000.00/ha (Table 4). The most substantial expenditure was land rental, comprising nearly half of the total cost, followed by key inputs such as seedlings, labor, fertilizers, pesticides, and organic manure. These factors collectively shaped the production expenses, which remained manageable and suggested the cultivar suitability for efficient cultivation. Over 2-year cultivation cycle, 'Nimas Agrihorti' achieved a cumulative yield of 12,000 kg, reflecting increasing trend an in productivity. The first year produced 3,000 kg with a revenue of IDR 30,000,000.00, while the second year yielded 9,000 kg, 90,000,000.00. generating IDR This upward trend indicates the plant's enhanced productivity as it matures, emphasizing its performance under optimal agronomic practices.

With total revenue reaching IDR 120,000,000.00 and net income at IDR 77,360,000.00, the farming of 'Nimas Agrihorti' proved to be highly profitable.

The R/C ratio of 2.81 further confirmed the economic sustainability of this cultivation. Efficient resource management enhanced the cost-benefit ratio, and the productivity gains in the second year reinforced the long-term potential of this citrus cultivar. Taken together, the agronomic strengths and economic returns position 'Nimas Agrihorti' as a promising commodity for both smallholder and commercial farming operations.

**Table 4.** Farm feasibility of *Citrusaurantifolia* 'Nimas Agrihorti'.

| No.   | Item       | Unit    | Quantity | Price      | Total (IDR) |
|-------|------------|---------|----------|------------|-------------|
|       |            |         |          | (IDR)      |             |
| 1     | Land rent  | ha      | 1        | 20,000,000 | 20,000,000  |
| 2     | Seeds      | Trees   | 600      | 15,000     | 9,000,000   |
| 3     | Manure     | Sacks   | 150      | 15,000     | 2,250,000   |
| 4     | NPK        | kg      | 200      | 12,000     | 2,400,000   |
|       | fertilizer |         |          |            |             |
| 5     | Pesticides | Package | 1        | 2,000,000  | 2,000,000   |
| 6     | Labour     | Person  | 2        | 50,000     | 7,000,000   |
|       | wages      | per day |          |            |             |
| Total | cost       |         |          |            | 42,640,000  |
| 1     | 'Nimas'    | kg      | 3,000    | 10,000     | 30,000,000  |
| 2     | 'Agrihorti | kg      | 9,000    | 10,000     | 90,000,000  |
|       | citrus'    |         |          |            |             |
| Total | revenue    |         |          |            | 120,000,000 |
| Total | income     |         |          |            | 77,360,000  |
| R/C   |            |         |          |            | 2.81        |

## Conclusion

'Nimas Agrihorti' is recognized as a promising citrus cultivar with significant potential development for as а biopharmaceutical product. Its morphological traits contributed to enhanced productivity under optimal agronomic conditions. Despite its relatively low sugar and acid levels, the fruit offered a mildly sweet flavor and contains a high concentration of vitamin C, supporting its classification as a source of bioactive compounds for therapeutic applications. The economic analysis reinforced its potential, with a favorable R/C ratio of 2.81, indicating high profitability. These attributes aligned well with the increasing biopharmaceutical demand for and functional food products in both domestic international and markets. Therefore, 'Nimas Agrihorti' represents a strong candidate for advancement as a medicinal high-value plant and agricultural commodity.

#### **Conflict of Interest**

All authors have no conflicts of interest to disclose.

#### Acknowledgement

We would like to extend our heartfelt gratitude to Sri Andayani, the technician of citrus germplasm, for her invaluable support and assistance throughout the research activities. Her expertise and dedication have greatly contributed to the success of this study.

#### References

- Adlini, M. N., Hafizah, D., & Umaroh, K. (2020). Karakterisasi tanaman jeruk (*Citrus* sp.) di kecamatan Nibung Hangus kabupaten Batu Bara Sumatera Utara (Characterization of citrus [*Citrus* sp.] plants in Nibung Hangus district, Batu Bara regency, North Sumatra). *KLOROFIL: Jurnal Ilmu Biologi dan Terapan*, 4(1), 1–7. https://doi.org/10.30821/kfl:jibt.v4i1.8921
- Alaqeel, N. K. (2024). Antioxidants from different citrus peels provide protection against cancer. *Brazilian Journal of Biology*, *84*. https://doi.org/10.1590/1519-6984. 271619
- Ali Abadi, S. M. H., Karami-Osboo, R., Kobarfard, F., Jahani, R., Nabi, M., Yazdanpanah, H., Mahboubi, A., Nasiri, A., & Faizi, M. (2022). Detection of lime juice adulteration by simultaneous determination of main organic acids using liquid chromatography-tandem mass spectrometry. *Journal of Food Composition and Analysis*, 105, 104223. https://doi.org/10.1016/ j.jfca.2021.104223
- Annisa, Z., Fareza, N. A., Cholissodin, I., & Muflikhah, L. (2022). Prediction of biopharmaceutical crop yields in Indonesia using the extreme learning machine method. Jurnal Pengembangan Teknologi Informasi dan Ilmu Komputer,

6(11), 5331–5338. https://doi.org/ 10.1051/bioconf/20249101016

- Anuchapreeda. S., Anzawa. R., Viriyaadhammaa, N., Neimkhum, W., Chaiyana, W., Okonogi, S., & Usuki, T. (2020). Isolation and biological activity of agrostophillinol from kaffir lime (Citrus hystrix) leaves. Bioorganic & Medicinal Chemistry Letters, 127256. https://doi.org/ 30(14), 10.1016/j.bmcl.2020.127256
- BPS-Statistics Indonesia (2024). *Produksi* tanaman buah-buahan, 2021–2023. Retrieved from https://www.bps.go.id/id/statisticstable/2/NjIjMg==/produksitanaman-buah-buahan.html
- BRIN-Indonesia National Research and Innovation Agency (2023). Discussing recent research in biopharmacy and pharmaceuticals, BRIN promotes a new platform for research collaboration. Retrieved from https://www.brin.go.id/news/ 116721
- Budiarto, K., Andrini, A., Budiyati, E., Mariana, B. D., Martasari, C., Mas'udah, S., Yulia, N. D., Ikarini, I., & Yulianti, F. (2024). Bioactive phytochemical contents on fruit peel of several citrus species. *BIO Web of Conferences*, *91*, 01016. https://doi.org/10.1051/bioconf/202 49101016
- CCRC UGM (2008). Key lime (Citrus aurantifolia). Faculty of Pharmacy, Universitas Gadjah Mada. Retrieved from https://ccrc.farmasi.ugm.ac.id/ensik lopedia/ensiklopedia-tanamanantikanker/j/jeruk-nipis
- da Silva, C. M. G., Contesini, F. J., Sawaya, A. C. H. F., Cabral, E. C., da Silva Cunha, I. B., Eberlin, M. N., & de Oliveira Carvalho, P. (2013). Enhancement of the antioxidant activity of orange and lime juices by flavonoid enzymatic deglycosylation. *Food Research*

*International*, *52*(1), 308–314. https://doi.org/10.1016/j.foodres.20 13.03.019

- Denaro, M., Smeriglio, A., & Trombetta, D. (2021). Antioxidant and antiinflammatory activity of citrus flavanones mix and its stability after *in vitro* simulated digestion. *Antioxidant*, *10*(140) 1–17. https://doi.org/10.1155/2021/24888 04
- de Luna, F. C. F., Ferreira, W. A. S., Casseb, S. M. M., & de Oliveira, E. H. C. (2023). Anticancer potential of flavonoids: an overview with an emphasis on tangeretin. *Pharmaceuticals*, *16*(9), 1229. https://doi.org/10.3390/ph1609122 9
- Ernawati, H. R, Suharjon, Fika, W., Sutopo, Rahmaniar, D., Husni, I., & Yunimar (2023). *Budidaya jeruk nipis Citrus aurantifolia*. Retrieved from https://hortikultura.pertanian.go.id/ wp-content/uploads/2024/11/ Budidaya-Jeruk-Nipis-Citrusaurantifolia\_watermark\_compresse d.pdf
- Fahrurin, W. A., Hadi, S., Sukarsosno, Susetyarini, R. E., & Permana, F. H. (2023). Kajian jenis-jenis tumbuhan berkhasiat obat yang dimanfaatkan untuk pengobatan oleh masyarakat kecamatan Sendang kabupaten Tulungagung (A study of medicinal plant species utilized for traditional healthcare by the community in Sendang district, Tulungagung regency). Jurnal Bioedukasi, 6(1). https://doi.org/10.1051/bioconf/202 49101016
- Gao, L., Gou, N., Amakye, W. K., Wu, J., & Ren, J. (2022). Bioactivity guided isolation and identification of phenolic compounds from *Citrus aurantium* L. with anti-colorectal cancer cells activity by UHPLC-Q-TOF/MS. *Current Research in Food Science*, 5, 2251–2260.

https://doi.org/10.1016/j.crfs.2022. 11.013

- Gebregziabher, B. S., Gebremeskel, H., Debesa, B., Ayalneh, D., Mitiku, T., Wendwessen, T., Habtemariam, E., Nur, S., & Getachew, T. (2023). Carotenoids: dietary sources, health functions, biofortification, marketing trend and affecting factors-a review. Journal of Agriculture and Food Research, 14, 100834. https://doi.org/10.1016/ j.jafr.2023.100834
- Hijaz, F., Al-Rimawi, F., Manthey, J. A., & Killiny, N. (2020). Phenolics, flavonoids and antioxidant capacities in *Citrus* species with different degree of tolerance to Huanglongbing. *Plant Signaling & Behavior*, 15(5), 1752447. https://doi.org/10.1080/15592324.2 020.1752447
- Ivakdalam, L. M., & Rehena, Z. (2020). Pengaruh rendaman jeruk nipis terhadap kandungan vitamin C dan pH minuman infused water. *Jurnal Agribisnis Perikanan*, *12*(2), 344– 349. https://doi.org/10.29239/ j.agrikan.12.2.344-349
- Khan, M. K., Zill-E-Huma, & Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33(1), 85–104. https://doi.org/10.1016/ j.jfca.2013.11.004
- Kumar, D., Ladaniya, M. S., Gurjar, M., Kumar, S., & Mendke, S. (2021). Quantification of flavonoids, phenols and antioxidant potential from dropped *Citrus reticulata* Blanco fruits influenced by drying techniques. *Molecules*, 26(14), 4159. https://doi.org/10.3390/ molecules26144159
- Li, G., Rouseff, R., Cheng, Y., Zhou, Q., & Wu, H. (2021). Comprehensive identification and distribution pattern of 37 oxygenated heterocyclic compounds in

commercially important citrus juices. LWT, 152, 112351. https://doi.org/10.1016/j.lwt.2021.1 12351

- Liu, S., Li, S., & Ho, C. T. (2022). Dietary bioactives and essential oils of lemon and lime fruits. *Food Science and Human Wellness*, *11*(4), 753– 764. https://doi.org/10.1016/ j.fshw. 2022.03.001
- Mitra, S., Lami, M. S., Uddin, T. M., Das, R., Islam, F., Anjum, J., Hossain, Md. J., & Emran, T. B. (2022). Prospective multifunctional roles and pharmacological potential of dietary flavonoid narirutin. *Biomedicine & Pharmacotherapy*, 150, 112932. https://doi.org/ 10.1016/j.biopha.2022.112932
- Mueed, A., Shibli, S., Al-Quwaie, D. A., Ashkan, M. F., Alharbi, M., Alanazi, H., Binothman, N., Aljadani, M., Majrashi, K. A., Huwaikem, M., Abourehab, M. A. S., Korma, S. A., & El-Saadony, M. T. (2023). Extraction. characterization of polyphenols from certain medicinal plants and evaluation of their antioxidant, antitumor, antidiabetic, antimicrobial properties, and potential use in human nutrition. Frontiers in Nutrition, 10, 1125106. https://doi.org/10.3389/fnut.2023.1 125106
- Pebrianto, E. O., Ikawati, & Putri, D. A. (2023). Feasibility analysis of patchouli farming business in Burana village, Tabulahan district, Mamasa regency. Indonesian Journal of Agricultural Research, 201–211. 5(03), https://doi.org/ 10.32734/injar.v5i03.9409
- Prommaban, A., & Chaiyana, W. (2022). Microemulsion of essential oils from citrus peels and leaves with anti-aging, whitening, and irritation reducing capacity. *Journal of Drug Delivery Science and Technology*, 69, 103188. https://doi.org/10.1016/ j.jddst.2022.103188

- Rahayu, R. S., Poerwanto, R., Efendi, D., & Widodo, W. D. (2020). Appropriate duration of drought stress for Madura tangerine flower induction. *Jurnal Hortikultura Indonesia*, *11*(2), 82–90. https://doi.org/ 10.29244/jhi.11.2.82-90
- Saini, R. K., Ranjit, A., Sharma, K., Prasad, P., Shang, X., Gowda, K. G. M., & Keum, Y. S. (2022). Bioactive compounds of citrus fruits: a review of composition and health benefits of carotenoids. flavonoids, limonoids, and terpenes. Antioxidants. 11(2), 239. https://doi.org/10.3390/antiox1102 0239
- Setyanti, Y. H., Anwar, S., & Slamet, W. (2013). Photosynthetic characteristics and phosphorus uptake of alfalfa (*Medicago sativa*) forage under different cutting heights and nitrogen fertilization levels. *Animal Agriculture Journal*, 2(1), 86–96. https://doi.org/ 10.3390/agriculture12081187
- Shafiee, S., & Minaei, S. (2018). Combined data mining/NIR spectroscopy for purity assessment of lime juice. *Infrared Physics & Technology*, 91, 193–199. https://doi.org/10.1016/ j.infrared.2018.04.012
- Shafreen, R. B., Lubinska-Szczygeł, M., Różańska, A., Dymerski, Т., Namieśnik, J., Katrich, E., & Gorinstein, S. (2018). Human serum interactions with phenolic and aroma substances of kaffir (Citrus hystrix) and key lime (Citrus aurantifolia) juices. Journal of Luminescence. 115-122. 201. https://doi.org/10.1016/j.jlumin.201 8.04.010

- Shimada, A., Ueno, H., Inagaki, M., and Yoshimitsu, H. (2022). Glutaminase inhibitory activity of umbelliferone isolated from kabosu (*Citrus* sphaerocarpa Hort. ex Tanaka). Natural Product Research, 36(2), 605–609. https://doi.org/10.1080/ 14786419.2020.1788553
- Tullio, V., Gasperi, V., Catani, M. V., & Savini, I. (2020). The impact of whole grain intake on gastrointestinal tumors: a focus on colorectal, gastric, and esophageal cancers. *Nutrients*, 13(1), 81. https://doi.org/:10.3390/nu1301008 1
- Yulianto, S., Jamilatun, M., & Pangesti, F. A. (2022). Analisis kadar vitamin C wedang jeruk nipis (Citrus aurantifolia) berdasarkan variasi suhu menggunakan metode spektrofotometri UV-Vis (Analysis of vitamin C content in lime drink [Citrus aurantifolia] based on temperature variations using UV-Vis spectrophotometry method). Jurnal Formil (Forum Ilmiah) Kesmas Respati, 7(2), 208–218. https://doi.org/10.35842/formil.v7i 2.441
- Yustiningsih, M. (2019). Light intensity and photosynthetic efficiency in shade plants and plants exposed to direct light. *Bioedu*, 4(2), 43–48. https://doi.org/10.32938/jbe.v4i2.3 85
- Zulkifli, Z., Mulyani, S., Saputra, R., & Pulungan, L. A. B. (2022). Relationship between the length and width of pineapple leaves and the quality of pineapple leaf fibers based on leaf position and duration of leaf soaking. *Jurnal Agrotek Tropika*, *10*(2), 247. https://doi.org/:10.23960/jat.v10i2. 5461